# Implementation with Interdependent Values\* PRELIMINARY AND INCOMPLETE

Richard McLean Department of Economics Rutgers University New Brunswick, NJ 08901 Andrew Postlewaite Department of Economics University of Pennsylvania Philadelphia, PA 19104

March 22, 2003

 $<sup>* {\</sup>rm Postlewaite}$  gratefully acknowledges support from the National Science Foundation. We thank Steve Matthews for helpful comments.

## 1. Introduction

There is a large literature aimed at characterizing the social choice functions that can be implemented in Bayes Nash equilibria. This literature typically takes agents' information as exogenously given and fixed throughout the analysis. While for some problems this may be appropriate, the assumption is problematic for others. A typical analysis, relying on the revelation principle, maximizes some objective function subject to truthful revelation being a Bayes equilibrium. It is often the case that truthful revelation is not "ex post incentive compatible", that is, for a given agent, there are some vectors of the other agents' types for which the agent may be better off by misreporting his type than truthfully revealing it. Truthful revelation, of course, may still be a Bayes equilibrium, because agents announce their types without knowing other agents types: choices must be made on the basis of their beliefs about other agents' types. The difficulty with assuming that agents' information is exogenous is that when truthful revelation is not ex post incentive compatible, agents have incentives to learn other agents' types. To the extent that an agent can, at some cost, learn something about other agents' types, agents' beliefs when a mechanism is applied must be treated as endogenous.

A planner who designs a mechanism for which truthful revelation is expost incentive compatible can legitimately ignore agents' incentives to engage in espionage to discover other agents' types, and consequently, expost incentive compatibility is desirable. The Clarke-Groves-Vickrey mechanism (hereafter  $CGV)^1$  for private values environments is a classic example of a mechanism for which truthful revelation is expost incentive compatible. For this mechanism, each agent submits his or her valuation for each possible choice. The mechanism selects the outcome that maximizes the sum of the agents' submitted valuations, and prescribes a transfer to each agent an amount equal to the sum of the values of the other agents for the outcome. With these transfers, each agent has a dominant strategy to reveal his valuation truthfully. Cremer and McLean (1985) (hereafter CM) consider a similar problem in which agents have private information, but interdependent valuations; that is, each agent's valuation can depend on other agents' information. They consider the mechanism design problem in which the aim is to maximize the revenue obtained from auctioning an object. They analyze revelation games in which agents announce their types, and construct transfers similar to those in the CGV mechanism. The transfers are such that for each outcome, (roughly) each agent receives a transfer equal to the sum of the valuations of the

<sup>&</sup>lt;sup>1</sup>See Clarke (1971), Groves (1973) and Vickrey (1961).

other agents. Because each agent's valuation depends on other agents' announced types, truthful revelation will not generally be a dominant strategy. Cremer and McLean show, however, that under certain conditions<sup>2</sup> truthful revelation will, as in the CGV mechanism, be expost incentive compatible.

There has recently been renewed interest in mechanisms for which truthful revelation is expost incentive compatible. Dasgupta and Maskin (2000), Perry and Reny (2002) and Ausubel (1999) (among others) have used the solution concept in designing auction mechanisms that assure an efficient outcome. Chung and Ely (2001) and Bergemann and Morris (2002) analyze the solution concept more generally. These papers (and Cremer and McLean), however, restrict attention to the case that agents' private information is one dimensional, a serious restriction for many problems. Consider, for example, a problem in which an oil field is to be auctioned, and each agent may have private information about the quantity of the oil in the field, the chemical characteristics of the oil, the capacity of his refinery to handle the oil and the demand for the refined products in his market, all of which affect this agent's valuation (and potentially other agents' valuations as well). While the assumption that information is single dimensional is restrictive. it is necessary: Jehiel et. al. (2002) show that for general mechanism design problems with interdependent values and multidimensional signals, for nearly all valuation functions, truthful revelation will be an expost equilibrium only for trivial outcome functions.

Thus, it is only in the case of single dimensional information that we can hope for ex post equilibria for interdependent value problems. But even in the single dimensional case, there are difficulties. Most work on mechanism design in problems with asymmetric information begin with utilities of the form  $u_i(c; t_i, t_{-i})$ , where c is a possible outcome,  $t_i$  represents agent i's private information and  $t_{-i}$  is a vector representing other agents' private information. In the standard interpretation,  $u_i$  is a reduced form utility function that gives agent i's utility of the outcome c under the particular circumstances likely to obtain given the agents' information. In the oil field problem above, for example, an agent's utility for the oil may depend on (among other things) the amount and chemical composition of the oil and the future demand oil products, and other agents' information affects i's (expected) value for the field insofar as i's beliefs about the quantity and composition of the oil and the demand for oil products are affected by their information. In this paper, we begin from this more primitive data in which i has a utility function  $v_i(\theta; t_i)$ , where  $\theta$  is a complete description of the state of

 $<sup>^{2}</sup>$ The conditions are discussed in section 3.

the world and  $t_i$  is his private information. For the oil example,  $\theta$  would include those things that affect *i*'s value for the oil – the amount and composition of the oil, the demand for oil, etc. The relationship between agents' private information and the state of the world is given by a probability distribution P over  $\Theta \times T$ . This formulation emphasizes the fact that other agents' information affects agent *i* precisely to the extent that it provides information about the state of the world.

The reduced form utility function that is normally the starting point for mechanism design analysis can be calculated from this more primitive structure:  $u(c,t) \equiv \Sigma_{\theta} v_i(\theta;t) P(\theta|t)$ . Most work that employs ex post incentive compatibility makes additional assumptions on the reduced form utility functions  $u_i$ . It is typically assumed that each agent's types are ordered, and that agents' valuations are monotonic in any agent's type. Further, it is assumed that the utility functions satisfy a single-crossing property: a movement of a given agent from one type to a higher type causes his valuation to increase at least as much as any other agent's valuation. We show that the conditions on the primitive data of the problem that would ensure that the reduced form utility functions satisfy the single crossing property are very stringent; the reduced form utility functions associated with very natural single dimensional information problems can fail to satisfy the single crossing property.

In summary, while expost incentive compatibility is desirable, nontrivial mechanisms for which truthful revelation is expost incentive compatible fail to exist for a large set of important problems. We introduce in this paper a notion of  $\varepsilon - ex$ post incentive compatibility: a mechanism is  $\varepsilon$ -ex post incentive compatibile if truthful revelation is expost incentive compatible with probability at least  $1 - \varepsilon$ . If truthful revelation is  $\varepsilon$ -ex post incentive compatible for a mechanism, agents' incentive to collect information about other agents' is bounded by  $\varepsilon$  times the maximal gain from espionage. If espionage is costly, a mechanism designer can be relatively comfortable in taking agents' beliefs as exogenous when  $\varepsilon$  is sufficiently small. We show that the existence of mechanisms for which there are  $\varepsilon$ -incentive compatible equilibria is related to the concept of informational size introduced in McLean and Postlewaite (2001, 2002). When agents have private information, the posterior probability distribution on the set of states of the world  $\Theta$  will vary depending on a given agent's type. Roughly, an agent's informational size corresponds to the maximal expected change in the posterior on  $\Theta$  as his type varies, fixing other agents' types. We show that for any  $\varepsilon$ , there exists  $\delta$  such that if each agent's informational size is less than  $\delta$ , there exists an efficient mechanism for which truthful revelation is an  $\varepsilon$ -incentive compatible equilibrium.

We describe the model in the next section and provide a brief history of expost incentive compatibility in Section 3. In Section 4 we introduce a generalized CGV mechanism, along with an alternative efficient mechanism.

#### 2. The Model

Let  $\Theta = \{\theta_1, .., \theta_m\}$  represent the finite set of states of nature and let  $T_i$  be the finite set of types of player i. Let C denote the set of social alternatives. Agent i's payoff is represented by a nonnegative function  $v_i : C \times \Theta \times T_i \to \Re_+$ . R: we should note that while this assumption by itself is innocuous, it will not be innocuous when combined with the assumption that there is a "status quo" that yields payoff 0. These together are what assure the possibility of efficient outcomes with no infusion of money, and together they rule out the standard problem of deciding whether or not to build a public good. We will assume that there exists  $c_0 \in C$  such that  $v_i(c_0, \theta, t_i) = 0$  for all  $(\theta, t_i) \in \Theta \times T_i$  and that there exists M > 0 such that  $v_i(\cdot, \cdot, \cdot) \leq M$  for each i. We will say that  $v_i$  satisfies the *pure common value property* if  $v_i$  depends only on  $(c, \theta) \in C \times \Theta$  and the *pure private value property* if  $v_i$  depends only on  $(c, t_i) \in C \times T_i$ .

Let  $(\tilde{\theta}, \tilde{t}_1, \tilde{t}_2, ..., \tilde{t}_n)$  be an (n+1)-dimensional random vector taking values in  $\Theta \times T$   $(T \equiv T_1 \times \cdots \times T_n)$  with associated distribution P where

$$P(\theta, t_1, ..., t_n) = Prob\{\theta = \theta, t_1 = t_1, ..., t_n = t_n\}.$$

We will make the following full support assumptions regarding the marginal distributions:  $P(\theta) = \operatorname{Prob}\{\tilde{\theta} = \theta\} > 0$  for each  $\theta \in \Theta$  and  $P(t_i) = \operatorname{Prob}\{\tilde{t}_i = t_i\} > 0$ for each  $t_i \in T_i$ . If X is a finite set, let  $\Delta_X$  denote the set of probability measures on X. The set of probability measures on  $\Theta \times T$  satisfying the full support conditions will be denoted  $\Delta_{\Theta \times T}^*$ 

In many problems with differential information, it is standard to assume that agents have utility functions  $u_i : C \times T \to R_+$  that depend on other agents' types. It is worthwhile noting that, while our formulation takes on a different form, it is equivalent. Given a problem as formulated in this paper, we can define  $u_i(c,t) = \sum_{\theta \in \Theta} [v_i(c, \theta, t_i)P(\theta|t)]$ . Alternatively, given utility functions  $u_i : C \times T \to R_+$ , we can define  $\Theta \equiv T$  and define  $v_i(c, t, t_i) = u_i(c, t)$ . Our formulation will be useful in that it highlights the nature of the interdependence: agents care about other agents' types to the extent that they provide additional information about the state  $\theta$ . A social choice problem is a collection  $(v_1, ..., v_n, P)$  where  $P \in \Delta_{\Theta \times T}^*$ . A social choice function is a mapping  $q: T \to C$  that specifies an outcome in C for each profile of announced types. We will assume that  $q(t) = c_0$  if  $t \notin T^*$ , where  $c_o$  can be interpreted as a status quo point. A mechanism is a collection  $\{q, x_i\}_{i \in N}$  where  $q: T \to C$  is a social choice function and the functions  $x_i: T \to \Re$  are transfer functions. For any profile of types  $t \in T^*$ , let

$$\hat{v}_i(c;t) = \hat{v}_i(c;t_{-i},t_i) = \sum_{\theta \in \Theta} v_i(c,\theta,t_i) P(\theta|t_{-i},t_i).$$

Although  $\hat{v}$  depends on P, we suppress this dependence for notational simplicity.

**Definition**: Let  $(v_1, ..., v_n, P)$  be a social choice problem. A social choice function is *efficient* if for each  $t \in T^*$ ,

$$q(t) \in \arg\max_{c \in C} \sum_{j \in N} \hat{v}_j(c; t).$$

**Definition**: Let  $(v_1, ..., v_n, P)$  be a social choice problem. A mechanism  $\{q, x_i\}_{i \in N}$  is:

strongly expost incentive compatible if truthful revelation is an expost dominant strategy equilibrium: for all i, all  $t_i, t'_i \in T_i$ , all  $\sigma_{-i} \in T_{-i}$  and all  $t_{-i} \in T_{-i}$ such that  $(t_{-i}, t_i) \in T^*$ ,

$$\hat{v}_i(q(\sigma_{-i}, t_i); t_{-i}, t_i) + x_i(\sigma_{-i}, t_i) \ge \hat{v}_i(q(\sigma_{-i}, t_i'); t_{-i}, t_i) + x_i(\sigma_{-i}, t_i').$$

ex post incentive compatible if truthful revelation is an ex post Nash equilibrium: for all i, all  $t_i, t'_i \in T_i$  and all  $t_{-i} \in T_{-i}$  such that  $(t_{-i}, t_i) \in T^*$ ,

$$\hat{v}_i(q(t_{-i}, t_i); t_{-i}, t_i) + x_i(t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t_i'); t_{-i}, t_i) + x_i(t_{-i}, t_i').$$

interim incentive compatible (IC) if for each  $i \in N$  and all  $t_i, t'_i \in T_i$ 

$$\sum_{\substack{(t_{-i},t_i)\in T^*\\(t_{-i},t_i)\in T^*}} \left[\hat{v}_i(q(t_{-i},t_i);t_{-i},t_i) + x_i(t_{-i},t_i)\right] P(t_{-i}|t_i)$$

$$\geq \sum_{\substack{(t_{-i},t_i)\in T^*\\(t_{-i},t_i)\in T^*}} \left[\hat{v}_i(q(t_{-i},t_i');t_{-i},t_i) + x_i(t_{-i},t_i')\right] P(t_{-i}|t_i)$$

ex post individually rational (XIR) if

 $\hat{v}_i(q(t);t) + x_i(t) \ge 0$  for all i and all  $t \in T^*$ .

feasible if for each  $t \in T^*$ ,

$$\sum_{j \in N} x_j(t) \le 0$$

balanced if for each  $t \in T^*$ ,

$$\sum_{j \in N} x_j(t) = 0$$

Clearly, strong ex post IC implies ex post IC which in turn implies interim IC. If  $\hat{v}_i(c;t)$  does not depend on  $t_{-i}$ , then the notions of ex post dominant strategy and ex post Nash equilibrium coincide.<sup>3</sup> We will need one more incentive compatibility concept.

**Definition**: Let  $\varepsilon \geq 0$ . A mechanism  $\{q, x_i\}_{i \in N}$  is  $\varepsilon$ - expost incentive compatible if for all i, all  $t_i, t'_i \in T_i$ ,

Pr 
$$ob\{(\tilde{t}_{-i}, t_i) \in T^* \text{ and } (\hat{v}_i(q(\tilde{t}_{-i}, t_i); \tilde{t}_{-i}, t_i) + x_i(\tilde{t}_{-i}, t_i))$$
  
 $-(\hat{v}_i(q(\tilde{t}_{-i}, t'_i); \tilde{t}_{-i}, t_i) + x_i(\tilde{t}_{-i}, t'_i)) \ge -\varepsilon |\tilde{t}_i = t_i\} \ge 1 - \varepsilon.$ 

Note that  $\{q, x_i\}_{i \in N}$  is a 0- expost incentive compatible mechanism if and only if  $\{q, x_i\}_{i \in N}$  is an expost incentive compatible mechanism.

## **3.** Historical Perspective

As mentioned in the introduction, the typical modeling approach to mechanism design with interdependent valuations begins with a collection of functions  $u_i$ :  $C \times T \to \Re$  as the primitive objects of study. In this approach, the elements of each  $T_i$  are ordered and a single crossing property (see below) is imposed. To our knowledge, the earliest construction of an expost IC mechanism in this framework appears in Cremer and McLean (1985). In their setup,  $T_i = \{1, 2, ..., m_i\}$  and

 $<sup>^{3}</sup>$ For a discussion of the relationship between ex post dominant strategy equilibrium, dominant strategy equilibrium, ex post Nash equilibrium and Bayes-Nash equilibrium, see Cremer and McLean (1985).

 $C = [0, \overline{c}]$  is an interval. Let  $u'_i(c, t_{-i}, t_i)$  denote the derivative of  $u_i(\cdot, t_{-i}, t_i)$  evaluated at  $c \in C$ .

**Definition**: Let q be a social choice rule. An E(xtraction)- mechanism is any mechanism  $\{q, x_i\}_{i \in N}$  satisfying

$$x_{i}(t_{-i}, t_{i}) = x_{i}(t_{-i}, 1) - \sum_{\sigma_{i}=2}^{t_{i}} \left[ u_{i}(q(t_{-i}, \sigma_{i}), t_{-i}, \sigma_{i}) - u_{i}(q(t_{-i}, \sigma_{i} - 1), t_{-i}, \sigma_{i}) \right]$$

whenever  $t_{-i} \in T_{-i}$  and  $t_i \in T_i \setminus \{1\}$ .

There are many E- mechanisms, depending on the choice of  $x_i(t_{-i}, 1)$  for each  $t_{-i} \in T_{-i}$ . In their 1985 paper, CM define such mechanisms and use them (in conjunction with a full rank condition) to derive their full extraction results. If q and  $u_i$  satisfy certain assumptions, then there exists an E-mechanism that implements q as an expost Nash equilibrium and is also expost individually rational. This is summarized in the next result.

**Theorem** 1: Suppose that

(i)

 $u'_i(c, t_{-i}, t_i + 1) \ge u'_i(c, t_{-i}, t_i) \ge 0$ 

for each  $i \in N, t_{-i} \in T_{-i}, t_i \in T_i \setminus \{m_i\}$  and  $c \in C$ . [This is assumption 2 in CM.]

(ii) The social choice rule q is monotonic in the sense that **FIX THIS** 

$$q(t_{-i}, t_i + 1) \ge q(t_{-i}, t_i)$$

for each  $i \in N, t_{-i} \in T_{-i}, t_i \in T_i \setminus \{m_i\}$ . Then any E-mechanism is expost IC. If, in addition,

$$u_i(0,t) = 0$$
 for all  $t \in T$ ,

then there exists an E-mechanism  $\{q, x_i\}_{i \in N}$  satisfying feasibility, ex post IC and ex post IR.

**Proof**: If assumptions (i) and (ii) are satisfied, then any E-mechanism is expost IC as a result of Lemma 2 in CM (1985). Suppose that, in addition,  $u_i(0,t) = 0$  for all  $t \in T$ . For each  $t_{-i}$ , define

$$x_i(t_{-i}, 1) = -u_i(q(t_{-i}, 1), t_{-i}, 1)$$

Feasibility follows from the assumption that  $u_i(q(t_{-i}, 1), t_{-i}, 1) \ge 0$  and the observation that  $u_i(q(t_{-i}, \sigma_i), t_{-i}, \sigma_i) - u_i(q(t_{-i}, \sigma_i - 1), t_{-i}, \sigma_i) \ge 0$  for each  $\sigma_i$ . Since the resulting E-mechanism is expost IC, it follows that

$$\begin{aligned} u_i(q(t_{-i}, t_i); t_{-i}, t_i) + x_i(t_{-i}, t_i) &\geq u_i(q(t_{-i}, 1); t_{-i}, t_i) + x_i(t_{-i}, 1) \\ &= \int_0^{q(t_{-i}, 1)} u_i'(y; t_{-i}, t_i) dy + x_i(t_{-i}, 1) \\ &\geq \int_0^{q(t_{-i}, 1)} u_i'(y; t_{-i}, 1) dy + x_i(t_{-i}, 1) \\ &= u_i(q(t_{-i}, 1); t_{-i}, 1) + x_i(t_{-i}, 1) \\ &= 0. \end{aligned}$$

It is important to point out that the family of E-mechanisms includes ex post IC mechanisms that are ex post IR but do not extract the full surplus (such as the mechanism defined in the proof of Theorem 1 above) as well as ex post IC mechanisms that extract the full surplus but are not ex post IR (such as the surplus extracting mechanisms constructed in CM (1985) that satisfy interim IR but not ex post IR.)

If one is interested in implementing a specific social choice rule (e.g., an expost efficient rule), then one must make further assumptions that guarantee that q satisfies the monotonicity condition (ii). This can be illustrated in the special case of a single object auction with interdependent valuations studied in CM (1985). In this case, a single object is to be allocated to one of n bidders. If i receives the object, his value is the nonnegative number  $w_i(t)$ . In this framework,  $q(t) = (q_1(t), ..., q_n(t))$  where each  $q_i(t) \ge 0$  and  $q_1(t) + \cdots + q_n(t) \le 1$  and

$$u_i(q(t_{-i}, t'_i); t_{-i}, t_i) + x_i(t_{-i}, t'_i) = q_i(t_{-i}, t'_i)w_i(t_{-i}, t_i) + x_i(t_{-i}, t'_i).$$

Finally, efficiency means that

$$\sum_{i \in N} q_i(t) w_i(t) = \max_{i \in N} \{ w_i(t) \}.$$

**Theorem 2**: Suppose that

(i) for each  $i \in N, t_{-i} \in T_{-i}, t_i \in T_i \setminus \{m_i\}$ 

$$w_i(t_{-i}, t_i) \le w_i(t_{-i}, t_i + 1)$$

(ii) For all 
$$i, j \in N, t_{-i} \in T_{-i}, t_i \in T_i \setminus \{m_i\}$$
  
 $w_i(t_{-i}, t_i) \ge w_j(t_{-i}, t_i) \Rightarrow w_i(t_{-i}, t_i + 1) \ge w_j(t_{-i}, t_i + 1)$   
 $w_i(t_{-i}, t_i) > w_j(t_{-i}, t_i) \Rightarrow w_i(t_{-i}, t_i + 1) > w_j(t_{-i}, t_i + 1)$ 

Then there exists an efficient, ex post IR, ex post IC auction mechanism.

Condition (ii) in Theorem 2 guarantees that i's probability of winning  $q_i(t_{-i}, t_i)$  is nondecreasing in i's type  $t_i$ . Other authors have employed a marginal condition that implies (ii) when bidders' values are drawn from an interval. Dasgupta and Maskin (2000) and Perry and Reny (2002) (in their work on ex post efficient auctions) and Ausubel (1999) (in his work on auction mechanisms) assume that types are drawn from an interval and that the valuation functions are differentiable and satisfy

(i')

$$\frac{\partial w_i}{\partial t_i}(t) \ge 0$$

and (ii')

$$\frac{\partial w_i}{\partial t_i}(t) \ge \frac{\partial w_j}{\partial t_i}(t).$$

These are the continuum analogues of the discrete assumptions in Theorem 2 above.

In this paper, we do not take the  $u_i : C \times T \to \Re$  as the primitive objects of study. Instead, we derive the reduced form  $\hat{v}_i : C \times T \to \Re$  from the valuation function  $v_i : C \times \Theta \times T_i \to R_+$  and the conditional distributions  $P_{\Theta}(\cdot|t)$ . In an auction framework (such as that studied in McLean and Postlewaite (2002)),

$$w_i(t) = \sum_{\theta} v_i(\theta, t_i) P_{\Theta}(\theta|t).$$

In this special case, the second condition is quite restrictive. For example, suppose that  $v_i(\theta, t_i) = \alpha_i \theta + \beta_i$  for each i where  $\alpha_i > 0$ . Then

$$w_i(t) = \alpha_i \sum_{\theta} \theta P_{\Theta}(\theta|t) + \beta_i := \alpha_i \overline{\theta}(t) + \beta_i.$$

Assuming that  $\overline{\theta}(\cdot)$  is differentiable, then the second condition (ii') is satisfied only if

$$(\alpha_i - \alpha_j) \frac{\partial \theta}{\partial t_i}(t) \ge 0$$

and

$$(\alpha_j - \alpha_i) \frac{\partial \overline{\theta}}{\partial t_j}(t) \ge 0$$

for each i and j. If it is also required that  $\frac{\partial w_i}{\partial t_i}(t) = \frac{\partial \overline{\theta}}{\partial t_i}(t) \ge 0$  and  $\frac{\partial w_j}{\partial t_j}(t) = \frac{\partial \overline{\theta}}{\partial t_j}(t) \ge 0$  with strict inequality for some t, then  $\alpha_i = \alpha_j$ . **TYPO** 

In this paper, we do not investigate the assumptions that  $v_i$  and  $P_{\Theta}(\cdot|t)$  would need to satisfy in order for Theorem 1 to be applicable to the reduced form  $\hat{v}_i$ . Indeed, we believe that such assumptions are prohibitively restrictive. Instead, we make certain assumptions regarding the distribution  $P \in \Delta_{\Theta \times T}^*$  but no assumptions regarding the primitive valuation function  $v_i$ .

## 4. A Generalized Clarke-Groves-Vickrey Mechanism

Let q be a social choice rule and define transfers as follows:

$$\begin{aligned} \alpha_i^q(t) &= \sum_{j \in N \setminus i} \hat{v}_j(q(t); t) - \max_{c \in C} \left[ \sum_{j \in N \setminus i} \hat{v}_j(c; t) \right] & \text{if } t \in T^* \\ &= 0 & \text{if } t \notin T^* \end{aligned}$$

The resulting mechanism  $(q, \alpha_i^q)$  is the generalized CGV mechanism with interdependent valuations (GCGV for short.) (Ausubel(1999) and Chung and Ely (2002) use the term generalized Vickrey mechanisms, but for a different class of mechanisms.) If  $\hat{v}_i$  depends only on  $t_i$  (as in the case when  $\tilde{\theta}$  and  $\tilde{t}$  are stochastically independent), then the GCGV mechanism reduces to the classical CGV mechanism and it is well known that, in this case, the CGV mechanism is ex post individually rational and satisfies strong ex post IC. It is straightforward to show that the GCGV mechanism is ex post individually rational and feasible. However, it will generally not even satisfy *interim* IC. First, we show that the GCGV mechanism is ex post IC when P satisfies a property called nonexclusive information.

**Definition:** A measure  $P \in \Delta_{\Theta \times T}^*$  satisfies nonexclusive information (NEI) if

$$t \in T^* \Rightarrow P_{\Theta}(\cdot|t) = P_{\Theta}(\cdot|t_{-i}) \text{ for all } i \in N.$$

**Proposition A:** Let  $\{v_1, .., v_n\}$  be a collection of payoff functions. If  $P \in \Delta^*_{\Theta \times T}$  exhibits nonexclusive information and if  $q: T \to C$  is an efficient social

choice rule for the problem  $\{v_1, ..., v_n, P\}$ , then the GCGV mechanism  $(q, \alpha_i^q)$  is expost IC and expost IR.

*Proof*: See appendix.

Nonexclusive information is a strong assumption. Our goal in this paper is to identify conditions under which we can modify the GCGV payments so that the new mechanism is interim IC and approximately ex post IC. In the next section we discuss the two main ingredients of our approximationn results: informational size and the variability of agents' beliefs.

# 5. Informational Size and Variability of Beliefs

#### 5.1. Informational Size

If  $t \in T^*$ , recall that  $P_{\Theta}(\cdot|t) \in \Delta_{\Theta}$  denotes the induced conditional probability measure on  $\Theta$ . A natural notion of an agent's informational size is the degree to which he can alter the best estimate of the state  $\theta$  when other agents are announcing truthfully. In our setup, that estimate is the conditional probability distribution on  $\Theta$  given a profile of types t. Any profile of agents' types t = $(t_{-i}, t_i) \in T^*$  induces a conditional distribution on  $\Theta$  and, if agent i unilaterally changes his announced type from  $t_i$  to  $t'_i$ , this conditional distribution will (in general) change. We consider agent i to be informationally small if, for each  $t_i$ , there is a "small" probability that he can induce a "large" change in the induced conditional distribution on  $\Theta$  by changing his announced type from  $t_i$  to some other  $t'_i$ . This is formalized in the following definition.

#### **Definition**: Let

$$I_{\varepsilon}^{i}(t_{i}',t_{i}) = \{t_{-i} \in T_{-i} | (t_{-i},t_{i}) \in T^{*}, (t_{-i},t_{i}') \in T^{*} \text{ and } ||P_{\Theta}(\cdot|t_{-i},t_{i}) - P_{\Theta}(\cdot|t_{-i},t_{i}')|| > \varepsilon\}$$

The *informational size* of agent i is defined as

$$\nu_i^P = \max_{t_i \in T_i} \max_{t_i' \in T_i} \min\{\varepsilon \ge 0 | \operatorname{Prob}\{\tilde{t}_{-i} \in I_{\varepsilon}^i(t_i', t_i) | \tilde{t}_i = t_i\} \le \varepsilon\}.$$

Loosely speaking, we will say that agent i is *informationally small* with respect to P if his informational size  $\nu_i^P$  is "small." If agent *i* receives signal  $t_i$  but reports  $t'_i \neq t_i$ , the effect of this misreport is a change in the conditional distribution on  $\Theta$ from  $P_{\Theta}(\cdot|t_{-i}, t_i)$  to  $P_{\Theta}(\cdot|t_{-i}, t'_i)$ . If  $t_{-i} \in I_{\varepsilon}(t'_i, t_i)$ , then this change is "large" in the sense that  $||P_{\Theta}(\cdot|\hat{t}_{-i}, t_i) - P_{\Theta}(\cdot|\hat{t}_{-i}, t'_i)|| > \varepsilon$ . Therefore,  $\operatorname{Prob}\{\tilde{t}_{-i} \in I_{\varepsilon}(t'_i, t_i)|\tilde{t}_i = t_i\}$ is the probability that i can have a "large" influence on the conditional distribution on  $\Theta$  by reporting  $t'_i$  instead of  $t_i$  when his observed signal is  $t_i$ . An agent is informationally small if for each of his possible types  $t_i$ , he assigns small probability to the event that he can have a "large" influence on the distribution  $P_{\Theta}(\cdot|t_{-i}, t_i)$ , given his observed type. Informational size is closely related to the notion of *nonexclusive information* studied in (Postlewaite and Schmeidler (1986)). If all agents have zero informational size, then P must satisfy NEI. In fact, we have the following easily demonstrated result:  $P \in \Delta^*_{\Theta \times T}$  satisfies NEI if and only if  $\nu^P_i = 0$  for each  $i \in N$ .

#### 5.2. Variability of Agents' Beliefs

Whether an agent *i* can be given incentives to reveal his information will depend on the magnitude of the difference between  $P_{T_{-i}}(\cdot|t_i)$  and  $P_{T_{-i}}(\cdot|t'_i)$ , the conditional distributions on  $T_{-i}$  given different types  $t_i$  and  $t'_i$  for agent *i*. To define the measure of variability, we first define a metric *d* on  $\Delta_{\Theta}$  as follows: for each  $\alpha, \beta \in \Delta_{\Theta}$ , let

$$d(\alpha,\beta) = \left\| \frac{\alpha}{||\alpha||_2} - \frac{\beta}{||\beta||_2} \right\|_2$$

where  $|| \cdot ||_2$  denotes the 2-norm. Hence,  $d(\alpha, \beta)$  measures the Euclidean distance between the Euclidean normalizations of  $\alpha$  and  $\beta$ . If  $P \in \Delta_{\Theta \times T}$ , let  $P_{\Theta}(\cdot|t_i) \in \Delta_{\Theta}$ be the conditional distribution on  $\Theta$  given that *i* receives signal  $t_i$  and define

$$\Lambda_i^P = \min_{t_i \in T_i} \min_{t_i' \in T_i \setminus t_i} d(P_{\Theta}(\cdot|t_i), P_{\Theta}(\cdot|t_i'))^2$$

This is the measure of the "variability" of the conditional distribution  $P_{\Theta}(\cdot|t_i)$  as a function of  $t_i$ .

As mentioned in the introduction, our work is related to that of Cremer and McLean (1985,1989). Those papers and subsequent work by McAfee and Reny (1992) demonstrated how one can use correlation to fully extract the surplus in certain mechanism design problems. The key ingredient there is the assumption that the collection of conditional distributions  $\{P_{T_{-i}}(\cdot|t_i)\}_{t_i\in T_i}$  is a linearly independent set for each i. This of course, implies that  $P_{T_{-i}}(\cdot|t_i) \neq P_{T_{-i}}(\cdot|t_i')$  if  $t_i \neq t_i'$ and, therefore, that  $\Lambda_i^P > 0$ . While linear independence implies that  $\Lambda_i^P > 0$ , the actual (positive) size of  $\Lambda_i^P$  is not relevant in the Cremer-McLean constructions, and full extraction will be possible. In the present work, we do not require that the collection  $\{P_{T_{-i}}(\cdot|t_i)\}_{t_i\in T_i}$  be linearly independent (or satisfy the weaker cone condition in Cremer and McLean (1988)). However, the "closeness" of the members of  $\{P_{T_{-i}}(\cdot|t_i)\}_{t_i\in T_i}$  is an important issue. It can be shown that for each *i*, there exists a collection of numbers  $\varsigma_i(t)$  satisfying  $0 \leq \zeta_i(t) \leq 1$  and

$$\sum_{t_{-i}} \left[ \varsigma_i(t_{-i}, t_i) - \varsigma_i(t_{-i}, t'_i) \right] P_{T_{-i}}(t_{-i}|t_i) > 0$$

for each  $t_i, t'_i \in T_i$  if and only if  $\Lambda_i^P > 0$ . This means that, if the posteriors  $\{P_{T_{-i}}(\cdot|t_i)\}_{t_i \in T_i}$  are all distinct, then the "incentive compatibility" inequalities above are strict. However, the expression on the left hand side decreases as  $\Lambda^P \to 0$ . Hence, the difference in the expected reward from a truthful report and from a false report will be very small if the conditional posteriors are very close to each other. Our results require that informational size and aggregate uncertainty be small relative to the variation in these posteriors.

# 6. Implementation and Informational Size

#### 6.1. The Results

Let  $\{z_i\}_{i\in N}$  be an *n*-tuple of functions  $z_i: T \to \Re_+$  each of which assigns to each  $t \in T$  a nonnegative number, interpreted as a "reward" to agent *i*. If  $\{q, x_i\}_{i\in N}$  is a mechanism, then the associated *augmented* mechanism is defined as  $\{q, x_i + z_i\}_{i\in N}$ .

**Theorem A**: Let  $(v_1, ..., v_n)$  be a collection of payoff functions.

(i) Suppose that  $P \in \Delta_{\Theta \times T}^*$  satisfies  $\Lambda_i^P > 0$  for each i and suppose that  $q: T \to C$  is an expost efficient social choice rule for the problem  $\{v_1, ..., v_n, P\}$ . Then there exists an augmented GCGV mechanism  $\{q, \alpha_i^q + z_i\}_{i \in N}$  for the social choice problem problem  $(v_1, ..., v_n, P)$  satisfying expost IR and interim IC.

(ii) For every  $\varepsilon > 0$ , there exists a  $\delta > 0$  such that, whenever  $P \in \Delta^*_{\Theta \times T}$  satisfies

$$\max_{i} \nu_i^P \le \delta \min_{i} \Lambda_i^P,$$

and whenever  $q: T \to C$  is an expost efficient social choice rule for the problem  $\{v_1, .., v_n, P\}$ , there exists an augmented GCGV mechanism  $\{q, \alpha_i^q + z_i\}_{i \in N}$  with  $0 \leq z_i(t) \leq \varepsilon$  for every *i* and *t* satisfying expost IR, interim IC and  $\varepsilon$ -expost IC.

#### 6.2. Discussion

Our results rely on the following key lemma.

**Lemma A:** Suppose that  $q: T \to C$  is an expost efficient social choice rule for the problem  $\{v_1, ..., v_n, P\}$ . If  $(t_{-i}, t_i), (t_{-i}, t'_i) \in T^*$ , then

$$\begin{aligned} & (\hat{v}_i(q(t_{-i}, t_i); t_{-i}, t_i) + \alpha_i^q(t_{-i}, t_i)) - (\hat{v}_i(q(t_{-i}, t_i'); t_{-i}, t_i) + \alpha_i^q(t_{-i}, t_i')) \\ \geq & -2M(n-1)||P_\Theta(\cdot|t_{-i}, t_i) - P_\Theta(\cdot|t_{-i}, t_i')|| \end{aligned}$$

In the case of the GCGV and egalitarian mechanisms, Lemma A provides an upper bound on the "ex post gain" to agent i when i's true type is  $t_i$  but i announces  $t'_i$  and others announce truthfully. If agents have zero informations size that is, if P exhibits nonexclusive information- then  $||P_{\Theta}(\cdot|t_{-i}, t_i) - P_{\Theta}(\cdot|t_{-i}, t'_i)|| =$ 0 if  $(t_{-i}, t_i), (t_{-i}, t'_i) \in T^*$ . Hence, truth is an ex post Nash equilibrium and Proposition A follows. If  $v_i$  does not depend on  $\theta$ , then (letting  $|\Theta| = 1$ ), we recover the classic dominant strategy result for the CGV mechanisms in the pure private values case.

If agent i is informationally small, then (informally) we can deduce from Lemma A that

$$\Pr{ob}\{||P_{\Theta}(\cdot|\tilde{t}_{-i},t_i) - P_{\Theta}(\cdot|\tilde{t}_{-i},t_i')|| \approx 0|\tilde{t}_i = t_i\} \approx 1$$

so truth is an "approximate" expost equilibrium for the CGCV in the sense that

$$\Pr{ob}\{(\hat{v}_i(q(t_{-i}, t_i); t_{-i}, t_i) + \alpha_i^q(t_{-i}, t_i)) - (\hat{v}_i(q(t_{-i}, t_i'); t_{-i}, t_i) + \alpha_i^q(t_{-i}, t_i')) \ge 0 | \tilde{t}_i = t_i\} \approx 1.$$

Lemma A has a second important consequence: if agent i is informationally small, then truth is an approximate Bayes-Nash equilibrium in the GCGV mechanism so the mechanism is *approximately* interim incentive compatible. More precisely, we can deduce from Lemma A that the interim expected gain from misreporting one's type is essentially bounded from above by one's informational size. If we want the mechanism to be *exactly* interim incentive compatible, then we must alter the mechanism (specifically, construct an augmented GCGV mechanism) in order to provide the correct incentives for truthful behavior. It is in this step that variability of beliefs plays a crucial role. To see this, first note that incentive compatibility of the augmented GCGV requires that

$$\sum_{\substack{(t_{-i},t_i)\in T^*\\:(t_{-i},t_i)\in T^*}} \left[ \left( \hat{v}_i(q(t_{-i},t_i);t_{-i},t_i) + \alpha_i^q(t_{-i},t_i) \right) - \left( \hat{v}_i(q(t_{-i},t_i');t_{-i},t_i) + \alpha_i^q(t_{-i},t_i') \right) \right] P(t_{-i}|t_i)$$

$$+ \sum_{i(t_{-i},t_i)\in T^*} \left( z_i(t_{-i},t_i) - z_i(t_{-i},t_i') \right) P(t_{-i}|t_i)$$

$$\geq 0$$

The first term is bounded from below by  $-K\nu_i^P$  where K is a positive constant independent of P. If  $\Lambda_i^P > 0$ , then there exists a collection of numbers  $\varsigma_i(t)$ satisfying  $0 \leq \zeta_i(t) \leq 1$  and

$$\sum_{t_{-i}} \left[ \varsigma_i(t_{-i}, t_i) - \varsigma_i(t_{-i}, t'_i) \right] P_{T_{-i}}(t_{-i}|t_i) > 0$$

for each  $t_i, t'_i \in T_i$ . By defining  $z_i(t_{-i}, t_i) = \eta \zeta_i(t_{-i}, t_i)$  and choosing  $\eta$  sufficiently large, then we will obtain interim incentive compatibility of the augmented GCGV mechanism. This is part (i) of Theorem A. As the informational size of an agent decreases, the minimal reward required to induce the truth also decreases. If  $\Lambda_i^P$ large enough relative to an agent's informational size  $\nu_i^P$ , then we can construct an augmented mechanism satisfying interim incentive compatibility. This is part (ii) of Theorem A.

## 7. Extensions:

#### 7.1. Lower bounded Mechanisms

In this section, we present a generalization of Theorem A based on an insight provided by Lemma A and we begin with some notation. In a typical implementation or mechanism design problem, one computes the mechanism for each instance of the data that defines the social choice problem. Therefore, in many if not most cases of interest, the mechanism is parametrized by the data defining the social choice problem. If we fix a profile  $(v_1, ..., v_n)$  of payoff functions, then we can analyze the parametric dependence of the mechanism on the probability distribution P and this dependence can be modelled as a mapping that associates a mechanism with each  $P \in \Delta_{\Theta \times T}^*$ . We will denote this mapping  $P \mapsto (q^P, x_1^P, ..., x_n^P)$ . For example, the mapping naturally associated with the GCGV mechanism is defined by

$$q^{P}(t) \in \arg \max_{c \in C} \sum_{j \in N} \sum_{\theta \in \Theta} v_{i}(c, \theta, t_{i}) P(\theta | t_{-i}, t_{i}) \text{ if } t \in T^{*}$$
$$q^{P}(t) = c_{0} \text{ if } t \notin T^{*}$$

and

$$\begin{aligned} x_i^P(t) &= \sum_{j \in N \setminus i} \sum_{\theta \in \Theta} v_i(q^P(t), \theta, t_i) P(\theta | t_{-i}, t_i) - \max_{c \in C} \left[ \sum_{j \in N \setminus i} \sum_{\theta \in \Theta} v_i(c, \theta, t_i) P(\theta | t_{-i}, t_i) \right] & \text{if } t \in T^* \\ &= 0 \text{ if } t \notin T^* \end{aligned}$$

# NOTE CHANGE IN DEF BELOW Rich: isn't a mechanism necessarily going to be lower bounded if we're in the finite case?

**Definition:** Let  $(v_1, ..., v_n)$  be a profile of payoff functions. For each  $P \in \Delta_{\Theta \times T}^*$ , let  $(q^P, x_1^P, ..., x_n^P)$  be a mechanism for the social choice problem  $(v_1, ..., v_n, P)$ . We will say that the mechanism  $(q^P, x_1^P, ..., x_n^P)$  is *lower bounded* if there exists a K(P) > 0 such that for all  $P \in \Delta_{\Theta \times T}^*$ ,

$$\left( \hat{v}_i(q^P(t_{-i}, t_i); t_{-i}, t_i) + x_i^P(t_{-i}, t_i) \right) - \left( \hat{v}_i(q^P(t_{-i}, t_i'); t_{-i}, t_i) + x_i^P(t_{-i}, t_i') \right)$$
  
 
$$\geq -K(P) ||P_{\Theta}(\cdot|t_{-i}, t_i) - P_{\Theta}(\cdot|t_{-i}, t_i')||$$

whenever  $(t_{-i}, t_i), (t_{-i}, t'_i) \in T^*$ . We will say that the mapping  $P \mapsto (q^P, x_1^P, ..., x_n^P)$  is *lower bounded* if for each  $P \in \Delta^*_{\Theta \times T}$ , the mechanism  $(q^P, x_1^P, ..., x_n^P)$  is lower bounded and the constant K(P) may be chosen independent of P.

If q is an efficient SCR, then Lemma A shows that the GCGV mechanism is lower bounded. From the definitions, it should also be clear that any expost IR, lower bounded mechanism will satisfy expost incentive compatibility if P exhibits nonexclusive information.

#### NOTE CHANGE IN STATEMENT BELOW

**Theorem B:** Let  $(v_1, ..., v_n)$  be a collection of payoff functions. For each  $P \in \Delta_{\Theta \times T}^*$ , let  $(q^P, x_1^P, ..., x_n^P)$  be an ex post IR mechanism for the SCP  $\{v_1, ..., v_n, P\}$  where  $q^P : T \to C$  is a social choice rule. Furthermore, suppose that  $P \mapsto (q^P, x_1^P, ..., x_n^P)$  is lower bounded. Then for every  $\varepsilon > 0$ , there exists a  $\delta > 0$  such that, whenever  $P \in \Delta_{\Theta \times T}^*$  satisfies

$$\max_{i} \nu_{i}^{P} \le \delta \min_{i} \Lambda_{i}^{P},$$

there exists an augmented mechanism  $\{q^P, x_i^P + z_i^P\}_{i \in \mathbb{N}}$  with  $0 \leq z_i^P(t) \leq \varepsilon$  for every *i* and *t* satisfying ex post IR, interim IC and  $\varepsilon$ -ex post IC.

#### 7.2. Example: Pure Common Values

The GCGV mechanism is lower bounded but a large class of lower bounded mechanisms are associated with efficient social choice rules generated by payoff functions satisfying the pure common value assumption.

**Lemma C**: Suppose that  $v_i : C \times \Theta \to \Re_+$  and suppose that  $q^P$  is an efficient social rule for the social choice problem  $\{v_1, ..., v_n, P\}$ . That is,

$$q^{P}(t) \in \arg\max_{c \in C} \sum_{j \in N} \sum_{\theta \in \Theta} v_{j}(c, \theta) P(\theta | t_{-i}, t_{i}).$$

For each P, let

$$\beta_i^P(t) = \frac{1}{n} \sum_{j \in N} \sum_{\theta \in \Theta} v_j(q^P(t), \theta) P(\theta|t) - \sum_{\theta \in \Theta} v_i(q^P(t), \theta) P(\theta|t).$$

Then  $P \mapsto (q^P, \beta_1^P, .., \beta_n^P)$  is lower bounded, balanced and ex post IR. Proof: See appendix

**Theorem C**: Let  $(v_1, ..., v_n)$  be a collection of payoff functions satisfying the pure common value assumption.

(i) Suppose that  $P \in \Delta_{\Theta \times T}^*$  satisfies  $\Lambda_i^P > 0$  for each i and suppose that  $q^P: T \to C$  is an expost efficient social choice rule for the problem  $\{v_1, .., v_n, P\}$ . Then there exists a mechanism  $\{q^P, x_i^P\}_{i \in N}$  for the social choice problem problem  $(v_1, .., v_n, P)$  satisfying expost IR and interim IC.

(ii) For every  $\varepsilon > 0$ , there exists a  $\delta > 0$  such that, whenever  $P \in \Delta^*_{\Theta \times T}$  satisfies

$$\max_{i} \nu_{i}^{P} \le \delta \min_{i} \Lambda_{i}^{P},$$

and whenever  $q^P : T \to C$  is an efficient social choice rule for the problem  $\{v_1, ..., v_n, P\}$ , there exists a mechanism  $\{q^P, \beta_i^P + z_i^P\}_{i \in N}$  for the social choice problem problem  $(v_1, ..., v_n, P)$  with  $0 \leq z_i^P(t) \leq \varepsilon$  for every *i* and *t* satisfying ex post IR, interim IC and  $\varepsilon$ -ex post IC. Furthermore,

$$0 \le \sum_{i \in N} \left( \beta_i^P + z_i^P \right) = \sum_{i \in N} z_i^P \le n\varepsilon.$$

#### 8. Asymptotic Results

In this section, we address the problem of implementation in the presence of many agents. In an appropriate replica framework (in particular, for conditionally independent sequences that we define below), agents will become informationally small as the number of agents grows. As a result, the rewards  $z_i$  that induce truthful behavior will also go to zero as the number of agents grows. It is also of interest to determine whether or not the total reward  $\sum_{i \in N} z_i^P$  goes to zero and we will study that question now.

#### 8.1. Notation and Definitions:

We will assume that all agents have the same finite signal set  $T_i = A$ . Recall that  $J_r = \{1, 2, ..., r\}$ . For each  $i \in J_r$ , let  $v_i^r : C \times \Theta \times A \to \Re_+$  denote the payoff to agent i. For any positive integer r, let  $T^r = A \times \cdots \times A$  denote the r-fold Cartesian product and let  $t^r = (t_1^r, .., t_r^r)$  denote a generic element of  $T^r$ .

#### 8.2. Replica Economies and the Replica Theorem

**Definition**: A sequence of prob measures  $\{P^r\}_{r=1}^{\infty}$  with  $P^r \in \Delta_{\Theta \times T^r}$  is a **conditionally independent sequence** if there exists  $P \in \Delta_{\Theta \times A}$  such that

(a) For each r and each  $(\theta, t_1, ..., t_r) \in \Theta \times T^r$ ,

$$P^{r}(t_{1}^{r},..,t_{r}^{r}|\theta) = \operatorname{Prob}\{\tilde{t}_{1}^{r} = t_{1}, \tilde{t}_{2}^{r} = t_{2},...,\tilde{t}_{r}^{r} = t_{r}|\tilde{\theta} = \theta\} = \prod_{i=1}^{r} P(t_{i}|\theta).$$

- (b) For every  $\theta$ ,  $\hat{\theta}$  with  $\theta \neq \hat{\theta}$ , there exists a  $t \in A$  such that  $P(t|\theta) \neq P(t|\hat{\theta})$ .
- (c) The marginal measure of  $P^2$  on  $T^2$  exhibits positive variability.

Because of the symmetry in the objects defining a conditionally independent sequence, it follows that, for fixed r, the informational size of each  $i \in J_r$  is the same. In the remainder of this section we will drop the subscript i and will write  $\nu^{P^r}$  for the value of the informational size of agents in  $J_r$ .

**Lemma D**: Suppose that  $\{P^r\}_{r=1}^{\infty}$  is a conditionally independent sequence. For every  $\varepsilon > 0$  and every positive integer k, there exists an  $\hat{r}$  such that

$$r^k \nu^{P^r} \le \varepsilon$$

whenever  $r > \hat{r}$ .

The proof is provided in the appendix and is an application of a large deviations result due to Hoeffding (1960). With this lemma, we can prove the following asymptotic result.

#### NOTE CHANGE IN STATEMENT BELOW

**Theorem D**: Suppose that  $\{P^r\}_{r=1}^{\infty}$  is a conditionally independent sequence. Let M and  $\varepsilon$  be positive numbers. Let  $\{(v_1^r, .., v_r^r)\}_{r\geq 1}$  be a sequence of payoff function profiles and for each r, let  $\{q^{P^r}(r), x_1^{P^r}(r), .., x_r^{P^r}(r)\}$  be an expost IR mechanism for the SCP  $(v_1^r, .., v_r^r, P^r)$ . Suppose that

(1)  $|v_i^r(\cdot, \cdot, \cdot)| \leq M$  for all r and  $i \in J_r$ 

(2) For each r,  $(q^{P^r}(r), x_1^{P^r}(r), ..., x_r^{P^r}(r))$  is lower bounded mechanism with constant  $K(P^r)$  and for some positive integer  $L, r^{-L}K(P^r) \to 0$  as  $r \to \infty$ .

Then there exists an  $\hat{r}$  such that for all  $r > \hat{r}$ , there exists an augmented mechanism  $(q^{P^r}(r), x_1^{P^r}(r) + z_1^r, ..., x_r^{P^r}(r) + z_r^r)$  for the social choice problem  $(v_1^r, ..., v_r^r, P^r)$  satisfying ex post IR and interim IC. Furthermore, for each  $t^r \in T^r$ ,  $z_i^r(t^r) \ge 0$  and  $\sum_{i \in J_r}^r z_i^r(t^r) \le \varepsilon$ .

**Corollary:** Suppose that  $\{P^r\}_{r=1}^{\infty}$  is a conditionally independent sequence. Let M and  $\varepsilon$  be positive numbers. Let  $\{(v_1^r, ..., v_r^r)\}_{r\geq 1}$  be a sequence of payoff function profiles and for each r, let  $\{q^{P^r}(r), \alpha_1^{P^r}(r), ..., \alpha_r^{P^r}(r)\}$  denote the GCGV mechanism for the SCP  $(v_1^r, ..., v_r^r, P^r)$ . Suppose that  $|v_i^r(\cdot, \cdot, \cdot)| \leq M$  for all r and  $i \in J_r$ .

Then there exists an  $\hat{r}$  such that for all  $r > \hat{r}$ , there exists an augmented GCGV mechanism  $(q^{P^r}(r), \alpha_1^{P^r}(r) + z_1^r, ..., \alpha_r^{P^r}(r) + z_r^r)$  for the social choice problem  $(v_1^r, ..., v_r^r, P^r)$  satisfying ex post IR and interim IC. Furthermore, for each  $t^r \in T^r$ ,  $z_i^r(t^r) \ge 0$  and  $\sum_{i \in J_r}^r z_i^r(t^r) \le \varepsilon$ .

#### 9. Discussion

1. Note that for the asymptotic results, the asymptotic revenue is full extraction from the highest value guy. This is because we extract all the surplus except the payments in the augmentation, and the augmentation payments go to zero. This plus the fact that the surplus the high value guy gets goes to zero since he's getting the object at the second highest value and the difference between the highest value and the second highest value goes to zero.

# 10. Proofs:

# 10.1. Proof of Lemma A:

First, consider the GCGV mechanism. Choose  $(t_{-i}, t_i), (t_{-i}, t'_i) \in T^*$ . Then

$$\hat{v}_i(q(t_{-i}, t_i); t_{-i}, t_i) + \alpha_i(t_{-i}, t_i) = \hat{v}_i(q(t_{-i}, t_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t_i); t_{-i}, t_i)$$
$$- \max_{c \in C} \left[ \sum_{j \in N \setminus i} \hat{v}_j(c; t_{-i}, t_i) \right]$$

and

$$\hat{v}_{i}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}) + \alpha_{i}(t_{-i}, t_{i}') = \hat{v}_{i}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}) + \sum_{j \in N \setminus i} \hat{v}_{j}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}) \\
- \sum_{j \in N \setminus i} \hat{v}_{j}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}) \\
+ \sum_{j \in N \setminus i} \hat{v}_{j}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}') - \max_{c \in C} \left[ \sum_{j \in N \setminus i} \hat{v}_{j}(c; t_{-i}, t_{i}') \right]$$

Since

$$\hat{v}_i(q(t_{-i}, t_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t_i); t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) \ge \hat{v}_i(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{-i}, t_i) + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t'_i); t_{$$

it follows that

$$\begin{aligned} & (\hat{v}_i(q(t_{-i}, t_i); t_{-i}, t_i) + x_i(t_{-i}, t_i)) - (\hat{v}_i(q(t_{-i}, t_i'); t_{-i}, t_i) + x_i(t_{-i}, t_i')) \\ & \geq \max_{c \in C} \left[ \sum_{j \in N \setminus i} \hat{v}_j(c; t_{-i}, t_i') \right] - \max_{c \in C} \left[ \sum_{j \in N \setminus i} \hat{v}_j(c; t_{-i}, t_i) \right] \\ & - \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t_i'); t_{-i}, t_i') + \sum_{j \in N \setminus i} \hat{v}_j(q(t_{-i}, t_i'); t_{-i}, t_i) \end{aligned}$$

Let

$$q^*(t_{-i}, t_i) \in \arg \max_{c \in C} \left[ \sum_{j \in N \setminus i} \hat{v}_j(c; t_{-i}, t_i) \right]$$

and let

$$q^*(t_{-i}, t'_i) \in \arg\max_{c \in C} \left[ \sum_{j \in N \setminus i} \hat{v}_j(c; t_{-i}, t'_i) \right].$$

Then

$$\begin{split} \max_{q \in C} \left[ \sum_{j \in N \setminus i} \hat{v}_j(q; t_{-i}, t'_i) \right] &- \max_{q \in C} \left[ \sum_{j \in N \setminus i} \hat{v}_j(q; t_{-i}, t_i) \right] \\ &= \left[ \sum_{j \in N \setminus i} \hat{v}_j(q^*(t_{-i}, t'_i); t_{-i}, t'_i) \right] - \left[ \sum_{j \in N \setminus i} \hat{v}_j(q^*(t_{-i}, t_i); t_{-i}, t_i) \right] \\ &= \left[ \sum_{j \in N \setminus i} \hat{v}_j(q^*(t_{-i}, t'_i); t_{-i}, t'_i) - \sum_{j \in N \setminus i} \hat{v}_j(q^*(t_{-i}, t_i); t_{-i}, t'_i) \right] \\ &+ \left[ \sum_{j \in N \setminus i} \hat{v}_j(q^*(t_{-i}, t_i); t_{-i}, t'_i) - \sum_{j \in N \setminus i} \hat{v}_j(q^*(t_{-i}, t_i); t_{-i}, t_i) \right] \\ &\geq \sum_{j \in N \setminus i} \hat{v}_j(q^*(t_{-i}, t_i); t_{-i}, t'_i) - \sum_{j \in N \setminus i} \hat{v}_j(q^*(t_{-i}, t_i); t_{-i}, t_i) \end{split}$$

Therefore,

$$\begin{aligned} &(\hat{v}_{i}(q(t_{-i},t_{i});t_{-i},t_{i})+x_{i}(t_{-i},t_{i}))-(\hat{v}_{i}(q(t_{-i},t_{i}');t_{-i},t_{i})+x_{i}(t_{-i},t_{i}'))\\ &\geq \sum_{j\in N\setminus i}\hat{v}_{j}(q^{*}(t_{-i},t_{i});t_{-i},t_{i}')-\sum_{j\in N\setminus i}\hat{v}_{j}(q^{*}(t_{-i},t_{i});t_{-i},t_{i})\\ &-\sum_{j\in N\setminus i}\hat{v}_{j}(q(t_{-i},t_{i}');t_{-i},t_{i}')+\sum_{j\in N\setminus i}\hat{v}_{j}(q(t_{-i},t_{i}');t_{-i},t_{i})\\ &\geq -2M(n-1)||P_{\Theta}(\cdot|t_{-i},t_{i})-P_{\Theta}(\cdot|t_{-i},t_{i}')||\end{aligned}$$

#### 10.2. Proof of Theorem A:

We prove part (ii) first. Choose  $\varepsilon > 0$ . Let

$$M = \max_{\theta} \max_{i} \max_{t_i} \max_{q \in C} v_i(q, \theta, t_i)$$

and let K be the cardinality of T. Choose  $\delta$  so that

$$0 < \delta < \frac{\varepsilon}{4M(n+1)\sqrt{K}}$$

Suppose that  $P \in \Delta^*_{\Theta \times T}$  satisfies

$$\max_{i} \nu_{i}^{P} \le \delta \min_{i} \Lambda_{i}^{P}.$$

Define  $\hat{\nu}^P = \max_i \nu_i^P$  and  $\Lambda^P = \min_i \Lambda_i^P$ . Therefore  $\hat{\nu}^P \leq \delta \Lambda^P$ . Since Now we define an augmented GCGV mechanism. For each  $t \in T$ , define

$$z_i(t_{-i}, t_i) = \varepsilon \frac{P_{T_{-i}}(t_{-i}|t_i)}{||P_{T_{-i}}(\cdot|t_i)||_2}$$

Since  $0 \leq \frac{P_{T_{-i}}(t_{-i}|t_i)}{||P_{T_{-i}}(\cdot|t_i)||_2} \leq 1$ , it follows that

$$0 \le z_i(t_{-i}, t_i) \le \varepsilon$$

for all  $i, t_{-i}$  and  $t_i$ .

The augmented CGV mechanism  $\{q, \alpha_i^q + z_i\}_{i \in N}$  is clearly expost efficient. Individual rationality follows from the observations that

$$\hat{v}_i(q(t);t) + \alpha_i^q(t) \ge 0$$

and

$$z_i(t) \ge 0.$$

Claim 1: Let K = |T|. Then

$$\sum_{(t_{-i},t_i)\in T^*} \left( z_i(t_{-i}|t_i) - z_i(t_{-i}|t_i') \right) P(t_{-i}|t_i) \ge \frac{\varepsilon}{2\sqrt{K}} \Lambda_i^P$$

**Proof of Claim 1**:

$$\sum_{(t_{-i},t_i)\in T^*} (z_i(t_{-i}|t_i) - z_i(t_{-i}|t_i')) P(t_{-i}|t_i) = \sum_{(t_{-i},t_i)\in T^*} (z_i(t_{-i}|t_i) - z_i(t_{-i}|t_i')) P(t_{-i}|t_i)$$

$$= \sum_{(t_{-i},t_i)\in T^*} \varepsilon \left[ \frac{P_{T_{-i}}(t_{-i}|t_i)}{||P_{T_{-i}}(\cdot|t_i)||_2} - \frac{P_{T_{-i}}(t_{-i}|t_i')}{||P_{T_{-i}}(\cdot|t_i')||_2} \right] P(t_{-i}|t_i)$$

$$= \frac{\varepsilon ||P_{T_{-i}}(\cdot|t_i)||_2}{2} \left\| \frac{P_{T_{-i}}(\cdot|t_i)}{||P_{T_{-i}}(\cdot|t_i)||_2} - \frac{P_{T_{-i}}(\cdot|t_i')}{||P_{T_{-i}}(\cdot|t_i')||_2} \right\|^2$$

$$\geq \frac{\varepsilon}{2\sqrt{K}} \Lambda_i^P$$

This completes the proof of Claim 1.

# Claim 2:

 $\sum_{(t_{-i},t_i)\in T^*} \left[ (\hat{v}_i(q(t_{-i},t_i);t_{-i},t_i) + x_i(t_{-i},t_i)) - (\hat{v}_i(q(t_{-i},t_i');t_{-i},t_i) + x_i(t_{-i},t_i')) \right] P(t_{-i}|t_i) \ge -5M\hat{\nu}^P$ 

# **Proof of Claim 2**: Define

$$A_{i}(t'_{i}, t_{i}) = \{t_{-i} \in T_{-i} | (t_{-i}, t_{i}) \in T^{*}, (t_{-i}, t'_{i}) \in T^{*}, ||P_{\Theta}(\cdot|t_{-i}, t_{i}) - P_{\Theta}(\cdot|t_{-i}t'_{i})|| > \hat{\nu}^{P}\}$$
  
and

$$B_{i}(t'_{i}, t_{i}) = \{t_{-i} \in T_{-i} | (t_{-i}, t_{i}) \in T^{*}, (t_{-i}, t'_{i}) \in T^{*}, ||P_{\Theta}(\cdot | t_{-i}, t_{i}) - P_{\Theta}(\cdot | t_{-i}t'_{i})|| \leq \hat{\nu}^{P}\}$$
  
and

$$C_i(t'_i, t_i) = \{ t_{-i} \in T_{-i} | (t_{-i}, t_i) \in T^*, (t_{-i}t'_i) \notin T^* \}$$

Since  $\nu_i^P \leq \hat{\nu}^P$ , we conclude that

$$\operatorname{Prob}\{\tilde{t}_{-i} \in A_i(t'_i, t_i) | \tilde{t}_i = t_i\} \le \nu_i^P \le \hat{\nu}^P.$$

In addition,

$$0 \le \hat{v}_i(q(t_{-i}, t_i); t_{-i}, t_i) + \alpha_i^q(t_{-i}, t_i) \le \hat{v}_i(q(t_{-i}, t_i); t_{-i}, t_i) \le M$$

for all  $i, t_i$  and  $t_{-i}$ . Therefore,

$$\begin{aligned} |\hat{v}_{i}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}) + \alpha_{i}^{q}(t_{-i}, t_{i}')| &= |\hat{v}_{i}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}) - \hat{v}_{i}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}') \\ &+ \hat{v}_{i}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}') + \alpha_{i}^{q}(t_{-i}, t_{i}')| \\ &\leq |\hat{v}_{i}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}) - \hat{v}_{i}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}')| \\ &+ |\hat{v}_{i}(q(t_{-i}, t_{i}'); t_{-i}, t_{i}') + \alpha_{i}^{q}(t_{-i}, t_{i}')| \\ &\leq 3M \end{aligned}$$

for all  $i, t_i, t_i'$  and  $t_{-i}$ . Applying the definitions, it follows that

$$\begin{split} &\sum_{t_{-i}\in A_{i}(t'_{i},t_{i})} \left[ (\hat{v}_{i}(q(t_{-i},t_{i});t_{-i},t_{i}) + \alpha_{i}^{q}(t_{-i},t_{i})) - (\hat{v}_{i}(q(t_{-i},t'_{i});t_{-i},t_{i}) + \alpha_{i}^{q}(t_{-i},t'_{i})) \right] P(t_{-i}|t_{i}) \\ &\geq -3M \sum_{t_{-i}\in A_{i}(t'_{i},t_{i})} P(t_{-i}|t_{i}) \\ &\geq -3M\hat{\nu}^{P}. \\ &\sum_{t_{-i}\in B_{i}(t'_{i},t_{i})} \left[ (\hat{v}_{i}(q(t_{-i},t_{i});t_{-i},t_{i}) + \alpha_{i}^{q}(t_{-i},t_{i})) - (\hat{v}_{i}(q(t_{-i},t'_{i});t_{-i},t_{i}) + \alpha_{i}^{q}(t_{-i},t'_{i})) \right] P(t_{-i}|t_{i}) \\ &\geq -2M(n-1) \sum_{t_{-i}\in B_{i}(t'_{i},t_{i})} ||P_{\Theta}(\cdot|t_{-i},t_{i}) - P_{\Theta}(\cdot|t_{-i}t'_{i})||P(t_{-i}|t_{i}) \\ &\geq -2M(n-1)\hat{\nu}^{P}. \end{split}$$

and

$$\begin{split} &\sum_{t_{-i}\in C_{i}(t'_{i},t_{i})} \left[ \left( \hat{v}_{i}(q(t_{-i},t_{i});t_{-i},t_{i}) + \alpha_{i}^{q}(t_{-i},t_{i}) \right) - \left( \hat{v}_{i}(q(t_{-i},t'_{i});t_{-i},t_{i}) + \alpha_{i}^{q}(t_{-i},t'_{i}) \right) \right] P(t_{-i}|t_{i}) \\ &= \sum_{t_{-i}\in C_{i}(t'_{i},t_{i})} \left[ \left( \hat{v}_{i}(q(t_{-i},t_{i});t_{-i},t_{i}) + \alpha_{i}^{q}(t_{-i},t_{i}) \right) - \left( \hat{v}_{i}(c_{0};t_{-i},t_{i}) + 0 \right) \right] P(t_{-i}|t_{i}) \\ &= \sum_{t_{-i}\in C_{i}(t'_{i},t_{i})} \left( \hat{v}_{i}(q(t_{-i},t_{i});t_{-i},t_{i}) + \alpha_{i}^{q}(t_{-i},t_{i}) \right) P(t_{-i}|t_{i}) \\ &\geq 0. \end{split}$$

Combining these observations completes the proof of the claim 2.

Applying Claims 1 and 2, it follows that

$$\begin{split} &\sum_{\substack{(t_{-i},t_i)\in T^* \\ (\hat{v}_i(q(t_{-i},t_i);t_{-i},t_i) + \alpha_i(t_{-i},t_i) + z_i(t_{-i},t_i)) P(t_{-i}|t_i) \\ &-\sum_{\substack{(t_{-i},t_i)\in T^* \\ (\hat{v}_i(q(t_{-i},t_i');t_{-i},t_i) + \alpha_i(t_{-i},t_i') + z_i(t_{-i},t_i')) P(t_{-i}|t_i) \\ &= \sum_{\substack{(t_{-i},t_i)\in T^* \\ (t_{-i},t_i)\in T^* \\ (z_i(t_{-i},t_i) - z_i(t_{-i},t_i')) P(t_{-i}|t_i) \\ &+ \sum_{\substack{(t_{-i},t_i)\in T^* \\ (z_i(T_{-i},t_i) - z_i(t_{-i},t_i')) P(t_{-i}|t_i) \\ &\geq \frac{\varepsilon}{2\sqrt{K}}\Lambda_i^P - 2(n+1)M\hat{\nu}^P \\ &\geq 0. \end{split}$$

and the proof of part (ii) is complete.

Part (i) follows from the computations in part (ii). We have shown that, for any positive number  $\alpha$ , there exists an augmented GCGV mechanism  $\{q, \alpha_i^q + z_i\}_{i \in N}$  satisfying

$$\sum_{\substack{(t_{-i},t_i)\in T^*\\ 2\sqrt{K}}} \left[ \left( \hat{v}_i(q(t_{-i},t_i);t_{-i},t_i) + \alpha_i^q(t_{-i},t_i) \right) - \left( \hat{v}_i(q(t_{-i},t_i');t_{-i},t_i) + \alpha_i^q(t_{-i},t_i') \right) \right] P(t_{-i}|t_i)$$

$$\geq \frac{\alpha}{2\sqrt{K}} \Lambda_i^P - 5M\hat{\nu}^P$$

for each i and each  $t_i, t'_i$ . If  $\Lambda^P_i > 0$  for each i, then  $\alpha$  can be chosen large enough so that incentive compatibility is satisfied. This completes the proof of part (i).

#### 10.3. Proof of Lemma D

Let  $P(\cdot|\theta)$  denote the conditional measure on A and we assume that  $P(\cdot|\theta) \neq P(\cdot|\hat{\theta})$ . Let  $t^r = (t_1^r, ..., t_r^r)$  so that  $\Pr ob\{\tilde{t}^r = t^r|\tilde{\theta} = \theta\} = P(t_1^r|\theta) \cdots P(t_r^r|\theta)$ . For each  $\alpha \in A$ , let  $f(t^r, \alpha) = \#\{i \leq r|t_i^r = \alpha\}$  and define  $f(t^r) = (f(t^r, \alpha))_{\alpha \in A}$ .

For each  $\theta$ , let

$$\rho(\theta) := \max_{\hat{\theta} \neq \theta} \prod_{\alpha \in A} \left[ \frac{P(\alpha|\hat{\theta})}{P(\alpha|\theta)} \right]^{P(\alpha|\theta)}$$

Using the same argument found in Gul and Postlewaite (see their equation 9) we deduce that  $\rho(\theta) < 1$ . It is easy to show (simply compute the logarithm) that

there exists a  $\delta > 0$  such that

$$\prod_{\alpha \in A} \left[ \frac{P_{\Theta}(\alpha|\hat{\theta})}{P_{\Theta}(\alpha|\theta)} \right]^{\frac{f(t^r|\alpha)}{r} - P(\alpha|\theta)} \le \frac{1}{\sqrt{\rho(\theta)}}$$

whenever  $\hat{\theta} \neq \theta$  and  $||\frac{f(t^r)}{r} - P(\cdot|\theta)|| < \delta$ . Letting  $R = \max_{\theta} \rho(\theta)$ , we conclude that  $||\frac{f(t^r)}{r} - P(\cdot|\theta)|| < \delta$  implies that

$$\frac{P_{\Theta}(\hat{\theta}|t^{r})}{P_{\Theta}(\theta|t^{r})} = \left[\prod_{\alpha \in A} \left[\frac{P_{\Theta}(\alpha|\hat{\theta})}{P_{\Theta}(\alpha|\theta)}\right]^{P(\alpha|\theta)} \prod_{\alpha \in A} \left[\frac{P_{\Theta}(\alpha|\hat{\theta})}{P_{\Theta}(\alpha|\theta)}\right]^{\frac{f(t^{r}|\alpha)}{r} - P(\alpha|\theta)}\right]^{r} \le \left[\rho(\theta)\frac{1}{\sqrt{\rho(\theta)}}\right]^{r} \le R^{r/2}$$

whenever  $\hat{\theta} \neq \theta$ . This in turn implies that

$$||\chi_{\theta} - P_{\Theta}(\cdot|t^{r})|| \le 2(m-1)R^{r/2}$$

where  $\chi_{\theta}$  is the Dirac measure with  $\chi_{\theta}(\theta) = 1$  and  $|\Theta| = m$ . To complete the argument, choose  $t_i, t'_i \in A$  and note that for all r sufficiently large,

$$\begin{aligned} \Pr{ob\{||P_{\Theta}(\cdot|\tilde{t}_{-i}^{r},t_{i}) - P_{\Theta}(\cdot|\tilde{t}_{-i}^{r},t_{i}^{r})||} &> 4(m-1)R^{r/2}|\tilde{\theta} = \theta\} \\ &\leq \Pr{ob\{\exists \alpha \in A : ||\chi_{\theta} - P_{\Theta}(\cdot|\tilde{t}_{-i}^{r},\alpha)|| > 2(m-1)R^{r/2}|\tilde{\theta} = \theta\}} \\ &\leq \Pr{ob\{\exists \alpha \in A : ||\frac{f(\tilde{t}_{-i}^{r},\alpha)}{r} - P_{\Theta}(\cdot|\theta)|| \ge \delta|\tilde{\theta} = \theta\}} \\ &\leq \Pr{ob\{||\frac{f(\tilde{t}^{r})}{r} - P_{\Theta}(\cdot|\theta)|| \ge \delta/2|\tilde{\theta} = \theta\}} \\ &\leq 2\exp(\frac{-r\delta^{2}}{2}) \end{aligned}$$

where the last inequality is due to Hoeffding (JASA, 1963). Hence, for all r sufficiently large,

$$\nu_i^P \le \max\{4(m-1)R^{r/2}, \frac{2\exp(\frac{-r\delta^2}{2})}{\beta}\}$$

where

$$\beta := \min_{\alpha \in A} P(\alpha).$$

# 10.4. Proof of Theoprem D

For each  $t \in T$ , define

$$z_i(t_{-i}, t_i) = \frac{\varepsilon}{r} \frac{P_{T_{i+1}}(t_{i+1}|t_i)}{||P_{T_{i+1}}(\cdot|t_i)||_2} \text{ if } i = 1, ..., r-1$$
$$= \frac{\varepsilon}{r} \frac{P_{T_1}(t_1|t_r)}{||P_{T_1}(\cdot|t_r)||_2} \text{ if } i = r$$

Since

$$0 \le z_i(t_{-i}, t_i) \le \frac{\varepsilon}{r}$$

for all  $i,\,t_{-i}$  and  $t_i$  so individual rationality of the augmented mechanism follows from the observations that

$$\hat{v}_i(q(t);t) + x_i(t) \ge 0$$

and

 $z_i(t) \ge 0.$ 

Claim 1: Let  $K = |T^2|$ . Then

$$\sum_{(t_{-i},t_i)\in T^*} \left( z_i(t_{-i}|t_i) - z_i(t_{-i}|t_i') \right) P(t_{-i}|t_i) \ge \frac{\varepsilon}{2\sqrt{K}} \Lambda_i^P$$

Proof of Claim 1:

$$\begin{split} \sum_{(t_{-i},t_{i})\in T^{r}} \left(z_{i}(t_{-i}|t_{i}) - z_{i}(t_{-i}|t_{i}')\right) P^{r}(t_{-i}|t_{i}) &= \sum_{(t_{-i},t_{i})\in T^{r}} \left(z_{i}(t_{-i}|t_{i}) - z_{i}(t_{-i}|t_{i}')\right) P^{r}(t_{-i}|t_{i}) \\ &= \sum_{(t_{-i},t_{i})\in T^{r}} \frac{\varepsilon}{r} \left[\frac{P_{T_{i+1}}(t_{i+1}|t_{i})}{||P_{T_{i+1}}(\cdot|t_{i})||_{2}} - \frac{P_{T_{i+1}}(t_{i+1}|t_{i}')}{||P_{T_{i+1}}(\cdot|t_{i})||_{2}}\right] P(t_{-i}|t_{i}) \\ &= \sum_{(t_{i+1},t_{i})\in T^{r}} \frac{\varepsilon}{r} \left[\frac{P_{T_{i+1}}(t_{i+1}|t_{i})}{||P_{T_{i+1}}(\cdot|t_{i})||_{2}} - \frac{P_{T_{i+1}}(t_{i+1}|t_{i}')}{||P_{T_{i+1}}(\cdot|t_{i})||_{2}}\right] P(t_{i+1}|t_{i}) \\ &\geq \frac{\varepsilon}{2r\sqrt{K}} \Lambda_{i}^{P^{2}} \end{split}$$

This completes the proof of Claim 1.

# Claim 2:

$$\sum_{(t_{-i},t_i)\in T^r} \left[ \left( \hat{v}_i(q(t_{-i},t_i);t_{-i},t_i) + x_i(t_{-i},t_i) \right) - \left( \hat{v}_i(q(t_{-i},t_i');t_{-i},t_i) + x_i(t_{-i},t_i') \right) \right] P^r(t_{-i}|t_i) \ge -5M\nu^{P^r}(t_{-i}|t_i) = -5M\nu^$$

**Proof of Claim 2**: Define

$$A_{i}(t'_{i}, t_{i}) = \{ t_{-i} \in T^{r}_{-i} | || P^{r}_{\Theta}(\cdot | t_{-i}, t_{i}) - P^{r}_{\Theta}(\cdot | t_{-i}t'_{i}) || > \hat{\nu}^{P^{r}} \}$$

and

$$B_{i}(t'_{i}, t_{i}) = \{t_{-i} \in T^{r}_{-i} | ||P^{r}_{\Theta}(\cdot|t_{-i}, t_{i}) - P^{r}_{\Theta}(\cdot|t_{-i}t'_{i})|| \le \hat{\nu}^{P^{r}}\}.$$

We conclude that

$$\operatorname{Prob}\{\tilde{t}_{-i} \in A_i(t'_i, t_i) | \tilde{t}_i = t_i\} \le \nu^{P^r}.$$

In addition,

$$0 \le \hat{v}_i^r(q(t_{-i}, t_i); t_{-i}, t_i) + x_i(t_{-i}, t_i) \le \hat{v}_i^r(q(t_{-i}, t_i); t_{-i}, t_i) \le M$$

for all  $i, t_i$  and  $t_{-i}$ . Therefore,

$$\begin{aligned} |\hat{v}_{i}^{r}(q(t_{-i},t_{i}');t_{-i},t_{i}) + x_{i}(t_{-i},t_{i}')| &= |\hat{v}_{i}^{r}(q(t_{-i},t_{i}');t_{-i},t_{i}) - \hat{v}_{i}^{r}(q(t_{-i},t_{i}');t_{-i},t_{i}') \\ &+ \hat{v}_{i}^{r}(q(t_{-i},t_{i}');t_{-i},t_{i}') + x_{i}(t_{-i},t_{i}')| \\ &\leq |\hat{v}_{i}^{r}(q(t_{-i},t_{i}');t_{-i},t_{i}) - \hat{v}_{i}^{r}(q(t_{-i},t_{i}');t_{-i},t_{i}')| \\ &+ |\hat{v}_{i}^{r}(q(t_{-i},t_{i}');t_{-i},t_{i}') + x_{i}(t_{-i},t_{i}')| \\ &\leq 3M \end{aligned}$$

for all  $i, t_i, t'_i$  and  $t_{-i}$ . Applying the definitions, it follows that

$$\begin{split} &\sum_{t_{-i}\in A_{i}(t'_{i},t_{i})} \left[ (\hat{v}_{i}(q(t_{-i},t_{i});t_{-i},t_{i}) + x_{i}(t_{-i},t_{i})) - (\hat{v}_{i}(q(t_{-i},t'_{i});t_{-i},t_{i}) + x_{i}(t_{-i},t'_{i})) \right] P^{r}(t_{-i}|t_{i}) \\ &\geq -3M \sum_{t_{-i}\in A_{i}(t'_{i},t_{i})} P^{r}(t_{-i}|t_{i}) \\ &\geq -3M\hat{\nu}^{P}. \\ &\sum_{t_{-i}\in B_{i}(t'_{i},t_{i})} \left[ (\hat{v}_{i}(q(t_{-i},t_{i});t_{-i},t_{i}) + x_{i}(t_{-i},t_{i})) - (\hat{v}_{i}(q(t_{-i},t'_{i});t_{-i},t_{i}) + x_{i}(t_{-i},t'_{i})) \right] P^{r}(t_{-i}|t_{i}) \\ &\geq -2MK^{r} \sum_{t_{-i}\in B_{i}(t'_{i},t_{i})} ||P^{r}_{\Theta}(\cdot|t_{-i},t_{i}) - P^{r}_{\Theta}(\cdot|t_{-i}t'_{i})||P^{r}(t_{-i}|t_{i}) \\ &\geq -2MK^{r}\nu^{P^{r}}. \end{split}$$

Combining these observations completes the proof of the claim 2.

Applying Claims 1 and 2, it follows that for sufficiently large r,

$$\begin{split} &\sum_{\substack{(t_{-i},t_i)\in T^r\\(\hat{v}_i(q(t_{-i},t_i);t_{-i},t_i) + \alpha_i(t_{-i},t_i) + z_i(t_{-i},t_i))} P^r(t_{-i}|t_i)} \\ &- \sum_{\substack{(t_{-i},t_i)\in T^r\\(\hat{v}_i(q(t_{-i},t_i');t_{-i},t_i) + \alpha_i^q(t_{-i},t_i')) - (\hat{v}_i(q(t_{-i},t_i');t_{-i},t_i) + \alpha_i^q(t_{-i},t_i'))]} P^r(t_{-i}|t_i)} \\ &= \sum_{\substack{(t_{-i},t_i)\in T^r\\(t_{-i},t_i)\in T^r}} \left[ \left( \hat{v}_i(q(t_{-i},t_i) - z_i(t_{-i},t_i')) P^r(t_{-i}|t_i) + \alpha_i^q(t_{-i},t_i') \right) P^r(t_{-i}|t_i) \right] P^r(t_{-i}|t_i) \\ &+ \sum_{\substack{(t_{-i},t_i)\in T^r\\(t_{-i},t_i)\in T^r}} \left( z_i(t_{-i},t_i) - z_i(t_{-i},t_i') \right) P^r(t_{-i}|t_i) \\ &\geq \frac{\varepsilon}{2r\sqrt{K}} \Lambda_i^{P^2} - 3M\nu^{P^r} - 2MK^r\nu^{P^r} \\ &= \frac{1}{r} \left[ \frac{\varepsilon}{2\sqrt{K}} \Lambda_i^{P^2} - 3Mr\nu^{P^r} - 2M\left(\frac{K^r}{r^L}\right) \left(r^{L+1}\nu^{P^r}\right) \right] \\ &\geq 0. \end{split}$$

and the proof is complete.

# 11. Bibliography

## References

- Ausubel, L. (1999), "A Generalized Vickrey Auction," mimeo, University of Maryland.
- [2] Bergemann, D. and S. Morris (2002), "Robust Mechanism Design," mimeo, Yale University.
- [3] Chung, K-S. and J. Ely (2001), "Ex-post Incentive Compatible Mechanism Design," mimeo, Northwestern University.
- [4] Clarke, E. (1971), "Multipart Pricing of Public goods," Public Choice 8, 19-33.
- [5] Cremer, J. and McLean, R. P. (1985), "Optimal Selling Strategies under Uncertainty for a Discriminatory Monopolist when Demands Are Interdependent," *Econometrica*, 53, 345-61.
- [6] Cremer, J. and McLean, R. P. (1988), "Full Extraction of the Surplus in Bayesian and Dominant Strategy Auctions," *Econometrica*, 56, 1247-57.
- [7] Dasgupta, P. S. and Maskin, E. (2000), "Efficient Auctions," Quarterly Journal of Economics 115, 341-388.
- [8] Groves, T. (1973), "Incentives in Teams," *Econometrica* 41, 617-631.
- [9] Jehiel, P., M. Meyer-ter-Vehn and B. Moldovanu(2002), "Potentials and Implementation: A Generic Impossibility Theorem," Working Paper, University College London and University of Mannheim.
- [10] Maskin, E. S. (1992), "Auctions and Privatization," In H. Siebert (ed.), Privatization, 115-36.
- [11] McAfee, P. and P. Reny (1992), "Correlated Information and Mechanism Design," *Econometrica* 60, 395-421.
- [12] McLean, R. and A. Postlewaite (2002), "Informational Size and Incentive Compatibility," *Econometrica* 70, 2421-2454.

- [13] McLean, R. and A. Postlewaite (2002), "Informational Size and Efficient Auctions," mimeo, University of Pennsylvania.
- [14] McLean, R. and A. Postlewaite (2001), "Efficient Auction Mechanisms with Interdependent Types and Multidimensional Signals," mimeo.
- [15] Milgrom, P. R. and Weber, R. J. (1982), "A Theory of Auctions and Competitive Bidding," *Econometrica*, 50, 1089-1122.
- [16] Perry, M. and Reny, P. J. (2002), "An Efficient Auction," *Econometrica* 70, 199-1213.
- [17] Postlewaite, A. and Schmeidler, D. (1986), "Implementation in Differential Information Economies," *Journal of Economic Theory*, June 1986, 39, 14-33.
- [18] Vickrey, W. (1961), "Counterspeculation, Auctions and Competitive Sealed tenders," *Journal of Finance* 16, 8-37.